

**MATRIX MODELS AND INTEGRABLE $C < 1$
OPEN STRING THEORIES***LAURENT HOUART[†]*Service de Physique Théorique, Université Libre de Bruxelles, Boulevard du Triomphe
CP 225, B-1050 Bruxelles, Belgium***ABSTRACT**

We study in the double scaling limit the two-matrix model which represents the sum over closed and open random surfaces coupled to an Ising model. The boundary conditions are characterized by the fact that the Ising spins sitting at the vertices of the boundaries are all in the same state. We obtain the string equation.

Since the discovery of the double scaling limit¹, important progress has been achieved in the understanding of open-closed string theories in (p, q) conformal minimal model backgrounds.

We have now a complete coherent description of the $(2m - 1, 2)$ conformal minimal models coupled to 2D gravity. These theories are described by the KdV hierarchy associated with the lie algebra $sl(2, C)$. Starting with the usual one-hermitian-matrix model¹, which describes the sum over closed surfaces, one finds that the KdV flow equations organize the operator structure of the $(2m - 1, 2)$ theories. Supplying “initial conditions” given by the string equation yields a complete description of closed string theories in the $(2m - 1, 2)$ backgrounds.

The operator content of the theories is fully understood. Indeed, the infinite number of operators found by BRST analysis in the Liouville description² is also present in the KdV formulation. The remaining KdV flows are identified studying the macroscopic loops^{3,4,5}. They correspond to boundary operators⁴ (the boundary length and the infinite number of operators associated with it). The KdV hierarchy associated to $sl(2, C)$ is thus the good framework to describe both closed and open string theories in $(2m - 1, 2)$ conformal minimal backgrounds.

Moreover, a direct study of the one-hermitian-matrix model supplemented with a logarithmic potential⁶—having the effect of adding surfaces with boundaries of finite extent in the partition function—has been performed in the double scaling limit⁷. This leads to the generalization of the previous string equation to the case of a

*Talk presented at the Journées Relativistes '93, Brussels, April, 1993.

[†]Aspirant FNRS, e-mail: lhouart@ulb.ac.be

non-zero open string coupling.

Finally, it has been shown⁸ that the $sl(2, C)$ mKdV models are a different description of the same open-closed string theories.

The description of the general (p, q) models^a coupled to 2D gravity is much more involved. Closed string theories in these backgrounds have been studied. They are realized in terms of multi-matrix models and are characterized in the double scaling limit by the generalized KdV hierarchies^{9,10,11}. For a given q the $sl(q, C)$ KdV hierarchy organizes the operator structure of the theory. Once more, the spectrum of the theory contains the operators found in the Liouville description but some remaining flows no longer admit an interpretation in terms of boundary operators⁴. Furthermore, the $sl(q, C)$ KdV hierarchy does not allow a description of the boundary length.

Related to this problem is the issue of finding a string equation describing open string theories in (p, q) backgrounds. In this case, when surfaces with boundaries of finite extent are added to the partition sum, one has to pay attention to the boundary problem. There are indeed $q - 1$ order parameters in the theory and the boundary conditions associated with them have to be fixed.

Recently a string equation for general open string theories in (p, q) backgrounds has been derived by Johnson¹² in the integrable model framework. This equation is obtained by studying the $sl(q, C)$ generalization of a mapping which transforms the $sl(2, C)$ τ -function characterizing the closed case into the $sl(2, C)$ τ -function of the open case. However by making use of the beautiful underlying mathematical structure and bypassing the matrix model route some physical informations are lost. It is very hard to know what are the boundary conditions associated with the string equation.

This talk is concerned with the boundary condition problem. Starting with the discrete formulation, I work out an explicit example: the open string theory in the $(4, 3)$ conformal background with fixed boundary conditions. In the spirit of ref.[7], using the orthogonal polynomial method, I perform the double scaling limit on the two-matrix model which represents the sum over closed and open random surfaces coupled to an Ising model (IM). The boundary conditions are characterized by the fact that the Ising spins sitting at the vertices of the boundaries are all in the same state. Here I present only the results and refer the reader to ref.[20] for more details.

The model is given by the following two-matrix model: $F = \ln Z$

$$Z = \int \mathcal{D}M_1 \mathcal{D}M_2 \exp\left(-\frac{N}{g} \text{Tr} V(M_1, M_2)\right) \quad (1)$$

where V is the asymmetrical potential:

$$V(M_1, M_2) = \frac{1}{2}M_1^2 - \frac{1}{4}M_1^4 + \frac{1}{2}M_2^2 - \frac{1}{4}M_2^4 - cM_1M_2 + \gamma \ln(1 - e^\mu M_1^2) \quad (2)$$

When $\gamma = 0$, F corresponds to the sum over closed surfaces coupled to an Ising model¹³. The two $N \times N$ hermitian matrices M_1, M_2 are used to represent the

^a We consider $p > q$.

two spin states. The parameter g is related to the bulk cosmological constant and N is the closed string coupling. The solution of the model in this case is well-known^{14,15,16}.

When $\gamma \neq 0$ the logarithmic potential has the effect of adding surfaces with arbitrary boundaries of finite extent to the partition function^{6,7}. In this case, F represents the sum over closed and open triangulated surfaces with the Ising spins sitting at the vertices of the boundaries all in the same state (e.g. all up). The parameter γ is the open string coupling. In the continuum limit μ will correspond to the parameter which couples to the boundary length as well as the parameter which couples to the boundary magnetization (there is no way to distinguish them in this model).

The model can be solved non-perturbatively using the well-known orthogonal polynomial method¹⁷. The string equation found in the double scaling limit is given by:

$$G^2 w'' - 3uw = 27\Gamma G R_0(z, M) \quad (3)$$

$$\begin{aligned} \frac{6}{5} \left[\frac{1}{3} u^3 + \frac{8}{(12)^2} G^4 u^{(4)} - \frac{1}{2} G^2 u'' u - \frac{1}{4} G^2 (u')^2 + w^2 \right] = z \\ + \frac{54}{5} G^2 \Gamma [2R_1(z, M) - R'_0(z, M)] \end{aligned} \quad (4)$$

where z is the continuum bulk cosmological constant, G is the renormalized closed string coupling, Γ is the renormalized open string coupling and $\frac{d^2}{dz^2} F = -\frac{1}{G^2} u(z)$. The Z_2 breaking field w is proportional to $\langle 1\sigma \rangle$. $R_0(z, M)$ is the resolvent and $R_1(z, M)$ is the first jet of the resolvent¹⁸ associated with the operator $L = G^3 \partial_z^3 - \frac{3}{4} G \{u, \partial\} + \frac{3}{2} w$, to wit:

$$R_0(z, M) \equiv \langle z \mid \frac{1}{M+L} \mid z \rangle \quad (5)$$

$$R_1(z, M) \equiv \partial_{\bar{z}} \langle z \mid \frac{1}{M+L} \mid \bar{z} \rangle \Big|_{\bar{z}=z} \quad (6)$$

The constant M , which comes from the bare parameter μ , is the renormalized boundary magnetization constant.

Now, using the Adler mapping¹⁸, we compute a system of two equations for the resolvent R_0 and the first jet R_1 . The result is:

$$\begin{aligned} -G^3 (2R_1 - R'_0)^{(3)} + \frac{3}{4} G u' (2R_1 - R'_0) + \frac{3}{2} G u (2R_1 - R'_0)' \\ = 3w' R_0 + \frac{9}{2} w R'_0 + 3M R'_0 \end{aligned} \quad (7)$$

$$\begin{aligned} \frac{1}{9} G^4 R_0^{(5)} - \frac{5}{6} G^2 u R_0^{(3)} - \frac{5}{4} G^2 u' R_0^{(2)} - \left(\frac{3}{4} G^2 u'' - u^2 \right) R'_0 - \left(\frac{1}{6} G^2 u^{(3)} - u u' \right) R_0 \\ = \frac{1}{2} G w' (2R_1 - R'_0) + \frac{3}{2} G w (2R_1 - R'_0)' + G M (2R_1 - R'_0)' \end{aligned} \quad (8)$$

We then multiply eq.(7) by $G(2R_1 - R'_0)$ and add to it eq.(8) multiplied by $3R_0$. It gives us a total differential which integrated once with respect to z leads to:

$$\begin{aligned} & \frac{3}{4}G^2uT^2 - G^4[TT^{(2)} - \frac{1}{2}(T')^2] + \frac{1}{3}G^4[R_0R_0^{(4)} - R'_0R_0^{(3)}] \\ & + \frac{1}{6}G^4(R''_0)^2 - \frac{5}{2}G^2uR_0R_0^{(2)} - \frac{5}{4}G^2[u'R_0R'_0 - u(R'_0)^2] \\ & - \frac{1}{2}G^2u^{(2)}R_0^2 + \frac{3}{2}u^2R_0^2 - 3GTR_0(M + \frac{3}{2}w) = \frac{2}{3}G^{-2} \end{aligned} \quad (9)$$

where $T \equiv 2R_1 - R'_0$.

We have fixed the integration constant using the fact that $R_0(u=w=0) = \frac{1}{3}G^{-1}M^{-\frac{2}{3}}$ and $R_1(u=w=0) = -\frac{1}{3}G^{-2}M^{-\frac{1}{3}}$.

The model (1) is thus completely solved in the double scaling limit. The solution is given by eqs.(3), (4) and (9). The results obtained here are compatible with those of ref.[12]. They confirm also that the $sl(3, C)$ $mKdV$ models are an equivalent description of open string theories in (p,3) backgrounds¹⁹.

References

1. E. Brézin and V. Kazakov, *Phys. Lett.* **B236** (1990) 144; M. Douglas and S. Shenker, *Nucl. Phys.* **B335** (1990) 635; D. Gross and A.A. Migdal, *Phys. Rev. Lett.* **64** (1990) 127; *Nucl. Phys.* **B340** (1990) 333.
2. B. Lian and G. Zuckerman *Phys. Lett.* **B254** (1991) 417.
3. T. Banks, M. Douglas, N. Seiberg and S. Shenker, *Phys. Lett.* **B238** (1990) 279.
4. E. Martinec, G. Moore and N. Seiberg, *Phys. Lett.* **B263** (1991) 190.
5. G. Moore, N. Seiberg and M. Staudacher, *Nucl. Phys.* **B362** (1991) 665.
6. V.A. Kazakov, *Phys. Lett.* **B237** (1990) 212.
7. I.K. Kostov, *Phys. Lett.* **B238** (1990) 181.
8. S. Dalley, C.V. Johnson, T.R. Morris and A. Watterstam, *Mod. Phys. Lett.* **A7** (1992) 2753.
9. M. Douglas, *Phys. Lett.* **B238** (1990) 176.
10. P. Ginsparg, M. Goulian, M.R. Plesser and J. Zinn-Justin, *Nucl. Phys.* **B342** (1990) 539.
11. P. Di Francesco and D. Kutasov, in *proceedings of the Cargese Workshop "Random Surfaces and Quantum Gravity"*, ed. O. Alvarez *et al.* (Plenum Press, New York 1991).
12. C.V. Johnson, Princeton preprint IASSNS-HEP-93/5 (hep-th 9301112).
13. V.A. Kazakov, *Phys. Lett.* **A119** (1986) 140; D.V. Boulatov and V.A. Kazakov, *Phys. Lett.* **B186** (1987) 379.
14. . Crnković, P. Ginsparg and G. Moore, *Phys. Lett.* **B237** (1990) 196.
15. E. Brézin, M. Douglas, V. Kazakov and S. Shenker, *Phys. Lett.* **B237** (1990) 43.
16. D. Gross and A.A. Migdal, *Phys. Rev. Lett.* **64** (1990) 717.
17. C. Itzykson and J.-B. Zuber, *J. Math. Phys.* **21** (1980) 411; D. Bessis, C. Itzykson and J.-B. Zuber, *Adv. Appl. Math.* **1** (1980) 109.
18. L.A. Dickey, *Soliton Equations and Hamiltonian Systems* (World Scientific, Singapore, 1991), pp. 17-20.
19. L. Houart, *Phys. Lett.* **B295** (1992) 37.
20. L. Houart, preprint ULB-TH-02/93 (hep-th 9303157), to appear in *Phys. Lett.* **B**.